







2023-2024 Formula COVID-19 Vaccines

Vaccine Type →		Pfizer			Moderna		Novavax	
Vaccine Information ↓		Comirnaty 12+ years	Infant/Toddler 6 months–4 years	Pediatric 5-11 years	Pediatric 6 months–11 years	Spikevax 12+ years	Adolescent/Adult 12+ years	
								
Packaging	Vaccine name, manufacturer	Comirnaty®, Pfizer	COVID-19 Vaccine, Pfizer	COVID-19 Vaccine, Pfizer	Covid-19 Vaccine, Moderna	Spikevax™, Moderna	COVID-19 Vaccine, Novavax	
	Packaging	Pre-filled Syringes	Yellow Cap, Multi-dose Vial	Blue Cap, Single-dose Vial	Dark Blue Cap, Single-dose Vial	Dark Blue Cap, Single-dose Vial	Royal Blue Cap, Multi-dose Vial	
	Doses Per Vial	1 dose	3 doses	1 dose	1 dose	1 dose	5 doses	
	Carton Size	10 doses	30 doses	10 doses	10 doses	10 doses	10 doses	
	NDC-Unit of Dose (box)	00069-2377-10	59267-4315-02	59267-4331-02	80777-0287-92	80777-0102-95	80631-0105-02	
	NDC-Unit of Dose (vial)	00069-2377-01	59267-4315-02	59267-4315-01	00069-2362-01	80777-0287-07	80777-0102-04	
	CVX Code	309	308	310	311	312	211	
	CPT Code	91320	91318	91319	91321	91322	91304	
Minimum Order	10 doses	30 doses	10 doses	10 doses	10 doses	10 doses		
Storage	Storage Limits Before Puncture:	Label vaccine with expiration and use-by dates. (applicable to all) →						
	ULT	⊘	Until expiration date at -90°C to -60°C (-130°F to -76°F)		⊘	⊘		
	Thermal Shipper	⊘	⊘			⊘	⊘	
	Freezer	⊘	⊘					⊘
	Refrigerator	Until expiration at 2-8°C (36-46°F)	Up to 10 weeks at 2° to 8°C (36°F to 46°F). Write the date on carton-not to exceed expiration.		Until expiration at -50°C to -15°C (-58°F to 5°F) Up to 30 days (not to exceed expiration date) at 2-8°C (36-46°F)		Until expiration at 2-8°C (36-46°F)	
	Expiration Date	As printed on carton and syringe	Check the label or Pfizer product website .		Check Moderna product website or QR code.		Check product website	
	Storage Limits After Puncture	Multi-dose vials only: Record puncture and use-by time on vial label.						
Use-By Limit (Discard Time After 1st Puncture)	N/A Discard after single use	Discard 12 hours after puncture. Keep at 2° to 25°C (36° to 77°F)	N/A Discard after single use	N/A. Discard after single use.		Discard 12 hours after puncture. Keep at 2° to 25°C (36° to 77°F)		
Administration	Diluent (supplied)	Do not dilute	1.1 mL per vial	Do not dilute	Do not dilute	Do not dilute	Do not dilute	
	Dose Volume & Dose	0.3 mL, 30 mcg dose	0.3 mL, 3 mcg dose	0.3 mL, 10 mcg dose	0.25 mL, 25 mcg **	0.5 mL, 50 mcg	0.5 mL, 5 mcg	
	Refrigerator Thaw Time (Do not refreeze)	N/A	2 hours in carton (2° to 8°C/36°F to 46°F)		45 minutes for single dose vial or 1 hour for syringe at 2°C to 8°C (36°F to 46°F). Let stand at room temp for 15 min before administering.		N/A	
	Room Temp Thaw Time (Do not refreeze)	N/A	Vial: 30 minutes at up to 25°C (77°F)		15 minutes for single dose vial at 15° to 25°C (59°)		N/A	
Total Time at Room Temp	4 hours	Up to 12 hours (including thaw time) at 8°C to 25°C		Store up to 24 hours at 8°C to 25°C (46°F to 77°F)		N/A		
Funding	Vaccines for Children (VFC) *	✓	✓	✓	✓	✓	✓	
	Bridge Access Program *	✓	⊘	⊘	⊘	✓	✓	
	Private *	✓	✓	✓	✓	✓	✓	

Funding Key

Vaccines for Children (VFC):

0 – 18 years of age AND

- Medicaid eligible
- Uninsured
- Underinsured and/or
- American Indian / Alaska Native

Bridge Access Program:

Uninsured or underinsured adults*
18 years of age and older**

*Adults with Medicaid or Medicare are not eligible for Bridge Access Program vaccine.

**Sites enrolled in both VFC and Bridge should use VFC vaccine to vaccinate uninsured and underinsured 18-year-olds.

Private:

Anyone of any age with private insurance.

- Children with CHIP insurance.
- Adults 19 and older with Medicaid or Medicare

* For patient eligibility, see the Funding Key to the right.

** Discard the vial and excess volume after extracting a single 0.25mL dose